



The use of Expert Judgements in quantifying uncertainties in risk assessment of food and feed safety

— Experiences and lessons learnt from the implementation
of EFSA's Expert Knowledge Elicitation Guidance —

Olaf Mosbach-Schulz (EFSA-AMU)
Andy Hart (FERA)

Delft, 05th July 17, COST


DISCLAIMER

The presentation was drafted under the sole responsibility of the authors and is not considered as an EFSA output.

The positions and opinions presented are those of the author alone and are not intended to represent the views of EFSA.


EFSA GUIDANCE ON EXPERT KNOWLEDGE ELICITATION

- Published 2014
- “A *systematic, documented and reviewable process* to retrieve expert judgements from a group of experts in the form of a probability distribution”
- Describes formal EFSA process for steering elicitation studies
- Contains standard protocols for Cooke, Sheffield and Delphi methods



EFSA DRAFT GUIDANCE ON UNCERTAINTY

- Published March 2016
- Trial period of 1 year
- When finalised, will be mandatory for all EFSA's scientific assessments
- Central role for expert judgement
- Adds more streamlined methods:
 - 'Semi-formal EKE'
 - 'Expert discussion'
 - Approximate probability scale



Ref: <https://www.efsa.europa.eu/en/topics/topic/uncertainty>
<https://www.efsa.europa.eu/sites/default/files/160321DraftGDUncertaintyInScientificAssessment.pdf>

BASIC PRINCIPLES OF UNCERTAINTY GUIDANCE

Mandatory:

- List identified uncertainties
- Characterise their combined impact on the assessment outcome
- Clear & unambiguous

Flexible:

- Choice of methods
- Degree of refinement
- Scalable to time and resources available
- **Fit for purpose**

Output required:

- Express in quantitative terms the combined effect of as many as possible of the identified sources of uncertainty
- ...in terms of the range & probabilities of possible outcomes
- ...and describe any uncertainties that cannot be quantified

COMBINED UNCERTAINTY

- We need to quantify the *combined* uncertainty
- *But* it is never possible to quantify *all* uncertainties individually
- Solution proposed in EFSA draft guidance:
 - quantify some uncertainties *individually*
 - quantify other uncertainties *collectively*
 - put them together to quantify *combined uncertainty*
 - *describe* any uncertainties that cannot be quantified

'STANDARDISED ASSESSMENT PROCEDURES'

- Common in many parts of EFSA's work – especially for assessment of regulated products
- Established in legislation or guidance documents
- Contain elements that are accepted by assessors and decision-makers as providing adequate cover for uncertainty
 - uncertainty factors, default values, conservative assumptions, etc.
- Uncertainty guidance approach:
 - Continue using existing procedures
 - Check for and assess 'non-standard' sources of uncertainty
 - Calibrate conservatism of the procedures when they come up for review

EXAMPLE - MYCOTOXINS IN FOOD

- EFSA Contaminants Panel (CONTAM Panel) opinion on T-2 and HT-2 toxins (EFSA, 2011) – *prior to draft uncertainty guidance*
- Standardised procedure for assess dietary risk to consumers
 - *Tolerable Daily Intake (TDI)* = lowest No Observed Adverse Effect Level from animal studies \div 100
 - *Exposure* = 95th percentile of specified population based on dietary surveys (consumption), concentration data and body weight
 - If Exposure \leq TDI then conclude 'no health concern'
- For T-2 and HT-2:
 - *Tolerable Daily Intake (TDI)* = 100 ng/kg bw/day
 - *Exposure*: 95th percentile for Toddlers (12-36 months)
 - 23 ng/kg bw/day assuming non-detects are zeroes
 - 91 ng/kg bw/day assuming non-detects = limit of detection
 - List of non-standard sources of uncertainty

MYCOTOXINS IN FOOD

Sources of uncertainty	Direction ^(a)
Uncertainty of the analytical measurements	+/-
Occurrence data on feed not representative for all feed materials in which T-2 and HT-2 toxins could be present	+/-
Effect of food and feed processing	+/-
High variability of feedstuffs used and feeding systems for livestock	+/-
Use of UB occurrence data in the exposure estimations	+
Use of LB occurrence data in the exposure estimations	-
Limited exposure data on infants	+/-
Limited data on exposures for vegetarians	+/-
No toxicokinetic data on T-2 and HT-2 toxins in humans and in most animal species	+/-
Lack of information on the contribution of the toxicity of HT-2 toxin and other metabolites to overall toxicity	+/-
Combined effects with other mycotoxins or other toxic substances in food and feed	+/-

(a): + = uncertainty with potential to cause over-estimation of exposure/risk; - = uncertainty with potential to cause under-estimation of exposure/risk

- Assessed collectively by expert judgement:**

The CONTAM Panel concluded that given the uncertainties, the risk assessment of human and animal exposure to the sum of T-2 and HT-2 toxins is more likely to over- than under-estimate the risk.

- Implied quantitative judgements:**

- *>50% probability that risk is over-estimated*
- *the magnitude of the individual uncertainties and how they combine*

MYCOTOXINS IN FOOD

- *Tolerable Daily Intake (TDI)* = 100 ng/kg bw/day
- *Exposure*: 95th percentile for Toddlers (12-36 months)
 - 23 ng/kg bw/day assuming non-detects are zeroes
 - 91 ng/kg bw/day assuming non-detects = limit of detection
- *Effect of other uncertainties*: >50% probability that risk is over-estimated
- **CONTAM Panel's overall conclusion is qualitative:**

Estimates of chronic dietary exposure for populations of all age groups to the sum of T-2 and HT-2 toxins based on the available occurrence data are below the group TDI of 100 ng/kg b.w., and therefore there is no health concern.

- 'No concern' implies judgements about (a) the probability of exceeding the TDI, and (b) whether this raises concern
- **Need to separate these judgements & make them transparent**

MAKING THE PROBABILITY JUDGEMENT TRANSPARENT

- *Tolerable Daily Intake (TDI)* = 100 ng/kg bw/day
- *Exposure*: 95th percentile for Toddlers (12-36 months)
 - 23 ng/kg bw/day assuming non-detects are zeroes
 - 91 ng/kg bw/day assuming non-detects = limit of detection
- *Implied measure of risk*: ratio of exposure to TDI
- Effect of other uncertainties: **>50% probability that risk is over-estimated**

>50% ? <50%



Probability that exposure
 exceeds the TDI – assess
 by *expert judgement*
 considering the identified
 sources of uncertainty

0.23

0.91

1 (exposure = TDI)

Ratio of Toddler exposure to TDI


Sources of uncertainty	Direction ^(a)
Uncertainty of the analytical measurements	+/-
Occurrence data on feed not representative for all feed materials in which T-2 and HT-2 toxins could be present	+/-
Effect of food and feed processing	+/-
High variability of feedstuffs used and feeding systems for livestock	+/-
Use of UB occurrence data in the exposure estimations	+
Use of LB occurrence data in the exposure estimations	-
Limited exposure data on infants	+/-
Limited data on exposures for vegetarians	+/-
No toxicokinetic data on T-2 and HT-2 toxins in humans and in most animal species	+/-
Lack of information on the contribution of the toxicity of HT-2 toxin and other metabolites to overall toxicity	+/-
Combined effects with other mycotoxins or other toxic substances in food and feed	+/-

(a) + = uncertainty with potential to cause over-estimation of exposure/risk; - = uncertainty with potential to cause under-estimation of exposure/risk

ELICITATION OF 'NON-STANDARD UNCERTAINTIES'

Sources of uncertainty	Direction ^(a)
Uncertainty of the analytical measurements	+/-
Occurrence data on feed not representative for all feed materials in which T-2 and HT-2 toxins could be present	+/-
Effect of food and feed processing	+/-
High variability of feedstuffs used and feeding systems for livestock	+/-
Use of UB occurrence data in the exposure estimations	+
Use of LB occurrence data in the exposure estimations	-
Limited exposure data on infants	+/-
Limited data on exposures for vegetarians	+/-
No toxicokinetic data on T-2 and HT-2 toxins in humans and in most animal species	+/-
Lack of information on the contribution of the toxicity of HT-2 toxin and other metabolites to overall toxicity	+/-
Combined effects with other mycotoxins or other toxic substances in food and feed	+/-

(a): + = uncertainty with potential to cause over-estimation of exposure/risk; - = uncertainty with potential to cause under-estimation of exposure/risk




- *What would Exposure/TDI be if the uncertainties were resolved?*
- Contextual considerations for choice of elicitation method:
 - Strong time constraint, few experts per topic (≥ 1)
 - Assessed by Working Group, adopted by Panel
 - Non-overlapping areas of expertise
 - Conclusions reached by consensus

ELICITATION OF 'NON-STANDARD UNCERTAINTIES'

Sources of uncertainty	Direction ^(a)
Uncertainty of the analytical measurements	+/-
Occurrence data on feed not representative for all feed materials in which T-2 and HT-2 toxins could be present	+/-
Effect of food and feed processing	+/-
High variability of feedstuffs used and feeding systems for livestock	+/-
Use of UB occurrence data in the exposure estimations	+
Use of LB occurrence data in the exposure estimations	-
Limited exposure data on infants	+/-
Limited data on exposures for vegetarians	+/-
No toxicokinetic data on T-2 and HT-2 toxins in humans and in most animal species	+/-
Lack of information on the contribution of the toxicity of HT-2 toxin and other metabolites to overall toxicity	+/-
Combined effects with other mycotoxins or other toxic substances in food and feed	+/-

(a): + = uncertainty with potential to cause over-estimation of exposure/risk; - = uncertainty with potential to cause under-estimation of exposure/risk

Probability term	Subjective probability range
Extremely likely	99-100%
Very likely	90-99%
Likely	66-90%
As likely as not	33-66%
Unlikely	10-33%
Very unlikely	1-10%
Extremely unlikely	0-1%



■ Approaches explored in Trial Period:

- 'Semi-formal' versions of Sheffield method (behavioural aggregation)
- Elicit separate distributions for exposure and TDI
- ...or elicit one distribution for ratio of exposure to TDI
- ...or elicit probability for exposure > TDI
 - using 'approximate probability scale' adapted from IPCC

RESISTANCE TO QUANTIFYING UNCERTAINTY

- 'There are too many uncertainties'
- 'I cannot quantify them'
- 'I can quantify some, but not my model uncertainty'
- 'It would be subjective and that's unscientific'
- 'Uncertainty is unquantifiable by definition' (after Knight)
- 'You cannot quantify unknown unknowns'
- 'Quantifying uncertainty implies excessive precision'
- '...requires extensive data'
- '...requires complex computations'
- '...requires excessive time or resource'
- 'It takes too much time'
- 'The Panel won't accept it'
- 'Our job is to say whether it's safe, or not safe'
- 'Risk managers and the public don't want to know about uncertainty'
- 'Communicating uncertainty will undermine confidence'

SUGGESTIONS/ADVICE WELCOME...


- What elicitation methods to apply in such cases?
- Any options for weighting/calibration of experts?

MORE APPLICATIONS

Judgements on Statements (true/false)

A livestock fetus feels pain or other negative affect during stunning or bleeding of the dam.¹

1% to 33% likely



Ref: [For details please refer to](#)

¹ = EFSA AHAW Panel (EFSA Panel on Animal Health and Animal Welfare), 2017. Scientific Opinion on the animal welfare aspects in respect of the slaughter or killing of pregnant livestock animals (cattle, pigs, sheep, goats, horses). EFSA Journal 2017;15(5):4782, 96 pp


² = EFSA PLH Panel (EFSA Panel on Plant Health), 2016. Scientific opinion on the risk to plant health of *Ditylenchus destructor* for the EU territory. EFSA Journal 2016;14(12):4602, 124 pp.

³ = EFSA AHAW Panel (EFSA Panel on Animal Health and Animal Welfare), 2015. Scientific Opinion on welfare aspects of the use of perches for laying hens. EFSA Journal 2015;13(6):4131, 70 pp

Estimation of quantities (risks)

Number of seed potatoes infested with the potato rot nematode imported into EU in the next year.²


1.3 with 50% uncertainty interval from 0.4 to 5 tubers per year



Prioritise options (rankings)

Preference of laying hens to use a specific perch during nights. (out of 10 different perches)³

Preferred is a „single perch 50cm above the slatted area“ (78% D vs C)



LESSONS LEARNT

- Framing questions in conceptual models
- Listing evidence and uncertainties
- Enable probabilistic judgements
- Combining quantified uncertainties
- Priority setting between essentials and full assessment
- Combined judgement on unquantified uncertainties

FRAMING THE QUESTIONS IN CONCEPTUAL MODELS

- Judgements should be **„in principle“ observable** to the experts
- All terminology should be described **in an agreed glossary**
- The quantity to judge should be **an outcome of a „theoretical“ experiment**
- The total risk assessment should be described **in a conceptual model**

LISTING EVIDENCE AND UNCERTAINTIES

- **All evidence is listed and summarized**, e.g. data sources and stats
- **Surrogates**, e.g. read-across, are actively searched
- **Limitations of all sources are discussed**, listed and qualitatively described (uncertainty table with direction, magnitude of bias)
- **Combination of evidence** by expert elicitation

ENABLE PROBABILISTIC JUDGEMENTS

- **Adapt the quantification method** to the problem:
 - Quantile ranges for numbers
 - Subjective belief for statements
 - Rankings for priorities
- Discuss **numerical representation versus precision**
- Select and **train one method** per quantification
- Combine **individual judgements with reasoning**, even when the latter is imperfect
- **Provide graphical feedback** to the experts

COMBINING QUANTIFIED UNCERTAINTIES

- A **mathematical formulation** of the conceptual model is preferable
- A **stepwise interpretation** of the conceptual model is helpful:
 - E.g. to get European estimates:
 - 1st step: Individual observations
 - 2nd step: Country specific estimates
 - 3rd step: European estimates
- A **combined judgement** is always possible

FIT FOR PURPOSE MODELLING

- Model screening enables to **select influencing factors**
 - E.g. Fast track estimates

- **Sensitivity analysis** is an important feedback to understand uncertainties and mitigation measures (e.g. for decision making)

- **Tiered approaches** are helpful:
 - Full formal procedure
 - Semi-formal procedure
 - Essentials

UNQUANTIFIED UNCERTAINTIES

**Nevertheless not all uncertainties
are individually assessable!**

- List and discuss **all unquantified uncertainties**
- **Allow a correction of the final estimate**
in view of the additional uncertainties

THANKS FOR YOUR ATTENTION



(Detail of the Nordic Pavilion by Terike Haapoja at 55th Venice Biennale, 2013, photograph by Olaf Mosbach-Schulz)

Olaf Mosbach-Schulz

olaf.mosbach-schulz@efsa.europa.eu

European Food Safety Authority (EFSA)

Assessment and Methodological Support Unit (AMU)

Andy Hart

FERA Science