

## STATISTICS FOR STOCHASTIC PROCESSES - EXERCISE SHEET 1

Mastermath, Spring Semester 2024, Jakob Söhl (j.soehl@tudelft.nl)

**Ex. 1.1.** The process  $X_t = \sigma W_t + bt$  with  $\sigma, b \in \mathbb{R}$  and  $W$  Brownian motion is observed at the time points  $0 = t_{0,n} < t_{1,n} < \dots < t_{n,n} = T_n$ . Denote  $\Delta X_{i,n} = X_{t_{i,n}} - X_{t_{i-1,n}}$  and  $\Delta t_{i,n} = t_{i,n} - t_{i-1,n}$ .

- a) Compute the MLE  $\hat{\theta}_{\text{MLE}}$  for the parameter  $\theta = (b, \sigma^2)$  and find conditions such that the MLE is consistent, i.e.,  $\hat{\theta}_{\text{MLE}} \xrightarrow{d} \theta$ .
- b) Assume that  $b$  is known. Compute the Fisher information for the parameter  $\sigma^2$ .

## STATISTICS FOR STOCHASTIC PROCESSES - EXERCISE SHEET 2

Mastermath, Spring Semester 2024, Jakob Söhl (j.soehl@tudelft.nl)

**Ex. 2.1.** Consider the SDE  $dX_t = aX_t dt + \sigma dW_t$ ,  $t \geq 0$ ,  $X_0 = X^{(0)} \in L^2$ . Make the ansatz  $X_t(\omega) = C_t(\omega)e^{at}$ . Apply the Itô formula to  $C_t(\omega)$  and derive the solution of the SDE in this way. For  $a < 0$  show that  $X_t \xrightarrow{d} N(0, -\sigma^2/(2a))$  as  $t \rightarrow \infty$ . For  $a < 0$  find  $X^{(0)}$  so that the solution of the SDE is stationary.

**Ex. 2.2.** a) Use the Itô formula to show  $\int_0^t W_s dW_s = \frac{1}{2}(W_t^2 - t)$  for Brownian motion  $W$ .

b) Let  $\hat{a}_T$  be the MLE in the Ornstein–Uhlenbeck model with time-continuous observations  $(X_t)_{t \in [0, T]}$  and initial condition  $X^{(0)} = 0$ . Consider  $\hat{a}_T$  as  $T \rightarrow \infty$  under  $\mathbb{P}^0$ , i.e., with true parameter  $a = 0$ . Show that  $\hat{a}_T$  is consistent and that  $T\hat{a}_T$  converges in distribution. Show that the limit is not a centred normal distribution.

### STATISTICS FOR STOCHASTIC PROCESSES - EXERCISE SHEET 3

Mastermath, Spring Semester 2024, Jakob Söhl (j.soehl@tudelft.nl)

**Ex. 3.1.** Consider the regression model  $Y_i = f(i/n) + \epsilon_i$ ,  $i = 1, \dots, n$ , where  $\epsilon_i$  are i.i.d. errors with  $\mathbb{E}[\epsilon_i] = 0$  and  $\text{Var}(\epsilon_i) < \infty$ . Let  $f : [0, 1] \rightarrow \mathbb{R}$  be differentiable and  $\|f'\|_\infty \leq M$ . For  $x \in [0, 1]$  define the estimator  $\hat{f}_n(x, h)$  by

$$\hat{f}_n(x, h) = \frac{\sum_{i=1}^n Y_i \mathbb{1}_{[x-h, x+h]}(i/n)}{\sum_{i=1}^n \mathbb{1}_{[x-h, x+h]}(i/n)} \quad \text{for } \sum_{i=1}^n \mathbb{1}_{[x-h, x+h]}(i/n) \neq 0$$

and  $\hat{f}_n(x, h) = 0$  otherwise. Show that  $|\hat{f}_n(x, n^{-1/3}) - f(x)| = O_{\mathbb{P}}(n^{-1/3})$ .

**Ex. 3.2.** Let  $dX_t = b(t) dt + \frac{\sigma}{\sqrt{n}} dW_t$ ,  $t \in [0, 1]$ ,  $X_0 = 0$ , where  $\sigma > 0$ ,  $b : [0, 1] \rightarrow \mathbb{R}$  and  $W$  is Brownian motion. For time-continuous observations  $(X_t)_{t \in [0, 1]}$  we define the estimator

$$\hat{b}_n(x, h) = \frac{\int_0^1 \mathbb{1}_{[x-h, x+h]}(t) dX_t}{\int_0^1 \mathbb{1}_{[x-h, x+h]}(t) dt}.$$

Show that for  $\alpha$ -Hölder continuous functions  $b$  with  $\alpha \in (0, 1]$ , for  $h = n^{-1/(2\alpha+1)}$  and  $x \in [0, 1]$

$$|\hat{b}_n(x, h) - b(x)| = O_{\mathbb{P}}(n^{-\alpha/(2\alpha+1)}).$$

## STATISTICS FOR STOCHASTIC PROCESSES - EXERCISE SHEET 4

Mastermath, Spring Semester 2024, Jakob Söhl (j.soehl@tudelft.nl)

**Ex. 4.1.** Let  $b$ ,  $\sigma$  and  $1/\sigma$  be bounded. Moreover, let  $b$  be differentiable and let  $\sigma$  be twice differentiable. Let  $s(x) = \int_0^x \exp(-\int_0^y 2b(z)/\sigma(z)^2 dz) dy$  and assume  $s(x) \rightarrow \pm\infty$  as  $x \rightarrow \pm\infty$ . Let  $(X_t)_{t \geq 0}$  satisfy the SDE  $dX_t = b(X_t) dt + \sigma(X_t) dW_t$  with  $X_0 = X^{(0)}$ .

The adjoint of the infinitesimal generator  $A$  is given by  $A^*g = -(bg)' + \frac{1}{2}(\sigma^2 g)''$  for  $g \in C_0^2(\mathbb{R})$ . From the theory of semigroups it follows that if we can find any non-negative  $m \in C_0^2(\mathbb{R})$ ,  $m \not\equiv 0$ , which has finite integral and satisfies  $A^*m = 0$ , then  $m$  is up to normalising the density of a stationary distribution.

- a) Show that  $Y_t = s(X_t)$  satisfies  $dY_t = \tilde{\sigma}(Y_t) dW_t$  with  $\tilde{\sigma}(y) = s'(s^{-1}(y))\sigma(s^{-1}(y))$ .
- b) Let  $\tilde{A}^*$  be adjoint of the infinitesimal generator of the SDE in part a). Find a non-negative function  $m \not\equiv 0$  satisfying  $\tilde{A}^*m = 0$ .
- c) Assume  $\tilde{G} = \int_{-\infty}^{\infty} 1/\tilde{\sigma}(y)^2 dy < \infty$  with  $\tilde{\sigma}$  is as in part a). Let  $Y$  be a random variable having density  $1/(\tilde{G}\tilde{\sigma}(y)^2)$ . Determine the density of  $X = s^{-1}(Y)$  in terms of  $\sigma$  and  $b$ .

**Ex. 4.2.** Consider the SDE of the linear factor model  $dX_t = \vartheta b(X_t) dt + \sigma(X_t) dW_t$  for a given  $\vartheta > 0$ . Let  $b$  and  $\sigma$  be known, measurable and bounded and let  $\inf_{x \in \mathbb{R}} \sigma^2(x) \geq \underline{\sigma}^2 > 0$ . Let there be  $M, \gamma > 0$  such that  $\text{sign}(x) \frac{2b}{\sigma^2}(x) \leq -\gamma$  for all  $x$  with  $|x| \geq M$ . Let  $(X_t)_{t \in [0, T]}$  be time-continuous observations of a stationary solution  $X$  of the SDE. Let  $\mathbb{E}[b(X_0)^2/\sigma(X_0)^2] > 0$ . Show that

$$\left( \int_0^T \frac{b(X_t)}{\sigma(X_t)} dW_t \right) / \left( \int_0^T \frac{b(X_t)^2}{\sigma(X_t)^2} dt \right)$$

has positive denominator with probability tending to one and is of order  $O_{\mathbb{P}}(1/\sqrt{T})$ .

## STATISTICS FOR STOCHASTIC PROCESSES - EXERCISE SHEET 5

Mastermath, Spring Semester 2024, Jakob Söhl (j.soehl@tudelft.nl)

**Ex. 5.1.** Let  $b$  and  $\sigma$  be measurable and locally bounded and let  $\inf_{x \in \mathbb{R}} \sigma^2(x) \geq \underline{\sigma}^2 > 0$ . Let there be  $M, \gamma > 0$  such that  $\text{sign}(x)2b(x)/\sigma(x)^2 \leq -\gamma$  for all  $x$  with  $|x| \geq M$ . Let  $X$  be a stationary solution and  $\mu$  be an invariant measure of the SDE  $dX_t = b(X_t)dt + \sigma(X_t)dW_t$ . Define  $Af(x) := \frac{1}{2}\sigma^2(x)f''(x) + b(x)f'(x)$ . For  $f, g \in C^2(\mathbb{R})$  with compact support show that  $A$  is symmetric with respect to  $\mu$ , i.e.,

$$\int_{-\infty}^{\infty} Af(x)g(x) d\mu(x) = \int_{-\infty}^{\infty} f(x)Ag(x) d\mu(x).$$

**Ex. 5.2.** Consider the process  $X_t = x_0 + \int_0^t \sigma(s) dW_s$  for  $t \in [0, T]$ , where  $T > 0$  fixed,  $x_0 \in \mathbb{R}$ ,  $W$  is Brownian motion and  $\sigma : [0, T] \rightarrow \mathbb{R}$  is a bounded deterministic function. For  $n \geq 1$  the process  $(X_t)_{t \in [0, T]}$  is observed at the times  $0 = t_{0,n} < t_{1,n} < \dots < t_{n,n} = T$ . Denote  $\Delta X_{i,n} = X_{t_{i,n}} - X_{t_{i-1,n}}$ ,  $\Delta t_{i,n} = t_{i,n} - t_{i-1,n}$  and  $\Delta_n = \max_{1 \leq i \leq n} \Delta t_{i,n}$ . Let  $g : [0, T] \rightarrow \mathbb{R}$ ,  $\alpha \in (0, 1]$  and  $R > 0$  be such that  $|g(t) - g(s)| \leq R|t - s|^\alpha$  and  $|g(t)| \leq R$  for all  $s, t \in [0, T]$ . Consider the estimator  $\hat{\Lambda}_n(g) = \sum_{i=1}^n g(t_{i-1,n})(\Delta X_{i,n})^2$  for  $\Lambda(g) = \int_0^T g(s)\sigma(s)^2 ds$ .

- a) Define  $M_n := \sum_{i=1}^n g(t_{i-1,n}) \left( (\Delta X_{i,n})^2 - \int_{t_{i-1,n}}^{t_{i,n}} \sigma(s)^2 ds \right)$ . Show that there exists a constant  $C > 0$  depending only on  $R$  and  $T$  such that  $\mathbb{E}[M_n^2] \leq C \|\sigma^4\|_\infty \Delta_n$ .
- b) Show that there exists a constant  $D > 0$  depending only on  $R$  and  $T$  such that

$$\mathbb{E}[(\hat{\Lambda}_n(g) - \Lambda(g))^2] \leq D \|\sigma^4\|_\infty \max(\Delta_n, \Delta_n^{2\alpha}).$$

## STATISTICS FOR STOCHASTIC PROCESSES - EXERCISE SHEET 6

Mastermath, Spring Semester 2024, Jakob Söhl (j.soehl@tudelft.nl)

**Ex. 6.1.** Consider the setting of Ex. 1.1 with  $n \rightarrow \infty$ ,  $T_n \rightarrow \infty$  and both  $b \in \mathbb{R}$  and  $\sigma^2 > 0$  unknown. Construct confidence intervals  $B_n$  and  $S_n$  for  $b$  and  $\sigma^2$ , respectively, which are asymptotic of level  $\alpha \in (0, 1)$ , i.e.,

$$\begin{aligned}\mathbb{P}(b \in B_n) &\rightarrow 1 - \alpha \quad \text{as } n \rightarrow \infty, \\ \mathbb{P}(\sigma^2 \in S_n) &\rightarrow 1 - \alpha \quad \text{as } n \rightarrow \infty.\end{aligned}$$

**Ex. 6.2.** In the setting of Ex. 3.1 let  $x \in (0, 1)$ ,  $nh_n \rightarrow \infty$  and  $nh_n^3 \rightarrow 0$  as  $n \rightarrow \infty$ . Define  $\sigma^2 := \text{Var}(\epsilon_1)$ .

- a) Show that  $\sqrt{nh_n}(\hat{f}_n(x, h_n) - f(x)) \rightarrow N(0, \sigma^2/2)$  as  $n \rightarrow \infty$ .
- b) With  $\sigma^2$  known construct confidence intervals  $I_n$  for  $f(x)$  which are of asymptotic level  $\alpha \in (0, 1)$ , i.e.,

$$\mathbb{P}(f(x) \in I_n) \rightarrow 1 - \alpha \quad \text{as } n \rightarrow \infty.$$

## STATISTICS FOR STOCHASTIC PROCESSES - EXERCISE SHEET 7

Mastermath, Spring Semester 2024, Jakob Söhl (j.soehl@tudelft.nl)

**Ex. 7.1.** Let  $X$  be a  $d$ -dimensional Lévy process with Lévy measure  $\nu$ .

- a) Let  $\int_{\mathbb{R}^d} |x| \mathbb{1}_{\{|x|>1\}} d\nu(x) < \infty$ . Show that  $\mathbb{E}[X_t] = \gamma_1 t$  with  $\gamma_1 = \gamma + \int_{\mathbb{R}^d} x \mathbb{1}_{\{|x|>1\}} d\nu(x)$ .
- b) Let  $d = 1$  and  $\int_{\mathbb{R}} x^2 d\nu(x) < \infty$ . Show that  $\mathbb{E}[X_t] = \gamma_1 t$  for  $\gamma_1$  as in a) and  $\text{Var}(X_t) = (\sigma^2 + \tilde{\nu}(\mathbb{R}))t$  for  $d\tilde{\nu}(x) = x^2 d\nu(x)$ .

**Ex. 7.2.** Let  $\varphi$  be the characteristic function of a random variable  $X$  on  $\mathbb{R}$  and  $\varphi_n$  the empirical characteristic function of  $n$  i.i.d. samples of  $X$ . For complex-valued random variables  $Z_i$  we define  $\text{Cov}_{\mathbb{C}}(Z_1, Z_2) = \mathbb{E}[Z_1 \bar{Z}_2] - \mathbb{E}[Z_1] \overline{\mathbb{E}[Z_2]}$  and  $\text{Var}_{\mathbb{C}}(Z_1) = \mathbb{E}[|Z_1 - \mathbb{E}[Z_1]|^2]$ .

- a) Show that  $\mathbb{E}[\varphi_n(u)] = \varphi(u)$ ,  $\text{Cov}_{\mathbb{C}}(\varphi_n(u), \varphi_n(v)) = \frac{1}{n}(\varphi(u-v) - \varphi(u)\varphi(-v))$ ,  $\text{Var}_{\mathbb{C}}(\varphi_n(u)) = \frac{1}{n}(1 - |\varphi(u)|^2) \leq \frac{1}{n}$ .
- b) Define  $\mathcal{C}_n(u) := \sqrt{n}(\varphi_n(u) - \varphi(u))$  and conclude from a) that  $\mathbb{E}[\mathcal{C}_n(u) \overline{\mathcal{C}_n(v)}] = \varphi(u-v) - \varphi(u)\varphi(-v)$ . Show that in combination with  $\overline{\mathcal{C}_n(u)} = \mathcal{C}_n(-u)$  this completely determines the covariance structure of all finite-dimensional distributions, i.e., for all  $u_1, \dots, u_k$  the covariance of  $\text{Re}(\mathcal{C}_n(u_1)), \text{Im}(\mathcal{C}_n(u_1)), \dots, \text{Re}(\mathcal{C}_n(u_k)), \text{Im}(\mathcal{C}_n(u_k))$ .
- c) Define  $\Gamma$  to be the centred complex-valued Gaussian process with  $\Gamma(-u) = \overline{\Gamma(u)}$  and  $\text{Cov}_{\mathbb{C}}(\Gamma(u), \Gamma(v)) = \varphi(u-v) - \varphi(u)\varphi(-v)$ . Use the multivariate central limit theorem to show that  $\mathcal{C}_n \xrightarrow{fidi} \Gamma$ , i.e., all finite-dimensional distributions converge weakly.

## STATISTICS FOR STOCHASTIC PROCESSES - EXERCISE SHEET 8

Mastermath, Spring Semester 2024, Jakob Söhl (j.soehl@tudelft.nl)

**Ex. 8.1.** Let  $R > 8\sqrt{d}$  for  $d \in \mathbb{N}$ . Suppose that the empirical characteristic process  $\mathcal{C}_n$  satisfies uniformly for  $n \in \mathbb{N}$  and  $K \geq 2$

$$\mathbb{P}\left(\max_{u \in [-K, K]^d} |\mathcal{C}_n(u)| \geq R\sqrt{\log(nK^2)}\right) \leq C(\sqrt{n}K)^{(64d-R^2)/(64d+64)}.$$

Show that the empirical characteristic function converges uniformly on compact sets in  $L^p$ ,  $p \geq 1$ , to the true characteristic function with rate  $(\log(n)/n)^{1/2}$ .

**Ex. 8.2.** The following result is known as Hoeffding's inequality: If  $X_1, \dots, X_n$  are mean zero independent random variables taking values in  $[b_i, c_i]$  for constants  $b_i < c_i$ ,  $i = 1, \dots, n$ , respectively, then for  $u > 0$

$$\mathbb{P}\left(\sum_{i=1}^n X_i > u\right) \leq \exp\left(-\frac{2u^2}{\sum_{i=1}^n (c_i - b_i)^2}\right)$$

of which an obvious consequence is (why?)

$$\mathbb{P}\left(\left|\sum_{i=1}^n X_i\right| > u\right) \leq 2 \exp\left(-\frac{2u^2}{\sum_{i=1}^n (c_i - b_i)^2}\right).$$

Provide a proof of this inequality. [Hint: You may find it useful to first prove the auxiliary result  $\mathbb{E}[\exp(vX_i)] \leq \exp(v^2(c_i - b_i)^2/8)$  for  $v > 0$ , and then use Markov's inequality in conjunction with a bound for the moment generating function of  $v \sum X_i$ .]



## STATISTICS FOR STOCHASTIC PROCESSES - EXERCISE SHEET 9

Mastermath, Spring Semester 2024, Jakob Söhl (j.soehl@tudelft.nl)

**Ex. 9.1.** Let  $K$  be a bounded function on  $\mathbb{R}$  such that  $(1/A) \int_0^A K(x) dx \rightarrow 0$  as  $A \rightarrow \pm\infty$ .

- a) For  $f = c\mathbb{1}_{[\alpha, \beta]}$  with  $c \in \mathbb{R}$  and  $0 \leq \alpha < \beta < \infty$  show that  $\int_0^\infty f(x)K(\lambda x) dx \rightarrow 0$  as  $\lambda \rightarrow \pm\infty$ .
- b) For  $f \in L^1([0, \infty))$  prove that  $\int_0^\infty f(x)K(\lambda x) dx \rightarrow 0$  as  $\lambda \rightarrow \pm\infty$ . [Hint: You may use that for every  $\epsilon > 0$  there exists a finite linear combination  $g_\epsilon$  of functions as in a) such that  $\int_0^\infty |f(x) - g_\epsilon(x)| dx < \epsilon$ .]
- c) For  $f \in L^1(\mathbb{R})$  conclude that  $\int_{-\infty}^\infty f(x)K(\lambda x) dx \rightarrow 0$  as  $\lambda \rightarrow \pm\infty$ .
- d) Show the Riemann–Lebesgue lemma: If  $f \in L^1(\mathbb{R})$ , then  $\int_{\mathbb{R}} f(x)e^{i\lambda x} dx \rightarrow 0$  as  $\lambda \rightarrow \pm\infty$ .

**Ex. 9.2.** Let  $\tilde{w}^{U_n}(u) := (1/U_n)\tilde{w}(u/U_n)$ , where  $\tilde{w}$  is a continuous function, supported on  $[0, 1]$  with  $\tilde{w}(u) > 0$  for  $u \in (0, 1)$ . Consider the optimisation problem

$$(\sigma_n^2, \lambda_n) := \operatorname{argmin}_{(\sigma^2, \lambda)} \int_0^\infty \tilde{w}^{U_n}(u) (\operatorname{Re} \psi_n(u) + \sigma^2 u^2/2 + \lambda)^2 du.$$

- a) Show that it is solved by  $\sigma_n^2 = \int_0^\infty w_\sigma^{U_n}(u) \operatorname{Re} \psi_n(u) du$  and  $\lambda_n = \int_0^\infty w_\lambda^{U_n}(u) \operatorname{Re} \psi_n(u) du$  and derive the form of  $w_\sigma^{U_n}$  and  $w_\lambda^{U_n}$  in terms of  $\tilde{w}^{U_n}$ . Verify that  $w_\sigma^{U_n}(u) = U_n^{-3} w_\sigma^1(u/U_n)$  and  $w_\lambda^{U_n}(u) = U_n^{-1} w_\lambda^1(u/U_n)$ .
- b) Derive the identities  $\int_0^{U_n} (-u^2/2) w_\sigma^{U_n}(u) du = 1$ ,  $\int_0^{U_n} w_\sigma^{U_n}(u) du = 0$ ,  $\int_0^{U_n} (-1) w_\lambda^{U_n}(u) du = 1$  and  $\int_0^{U_n} (-u^2/2) w_\lambda^{U_n}(u) du = 0$ .

## STATISTICS FOR STOCHASTIC PROCESSES - EXERCISE SHEET 10

Mastermath, Spring Semester 2024, Jakob Söhl (j.soehl@tudelft.nl)

**Ex. 10.1.** Denote by  $\mathcal{F}f$  the Fourier transform of  $f$ . For an integer  $s \geq 0$  suppose that  $v$  satisfies  $\max_{k \in \{0, 1, \dots, s\}} \|v^{(k)}\|_{L^2(\mathbb{R})} \leq C$  and  $\|v^{(s)}\|_{\infty} \leq C$  for some  $C > 0$ , where  $v^{(k)}$  is the  $k$ -th derivative of  $v$ . Let  $w_{\gamma}$  and  $w_{\lambda}$  be functions supported on  $[0, 1]$  such that  $w_{\gamma}(\bullet)/(\bullet)^s$ ,  $w_{\lambda}(\bullet)/(\bullet)^s \in L^2(\mathbb{R})$  and  $\mathcal{F}[w_{\gamma}(\bullet)/(\bullet)^s], \mathcal{F}[w_{\lambda}(\bullet)/(\bullet)^s] \in L^1(\mathbb{R})$ . Define  $w_{\gamma}^{U_n}(u) = U_n^{-2}w_{\gamma}(u/U_n)$  and  $w_{\lambda}^{U_n}(u) = U_n^{-1}w_{\lambda}(u/U_n)$ . Show that

$$\left| \int_0^{\infty} w_{\gamma}^{U_n}(u) \operatorname{Im}(\mathcal{F}v(u)) \, du \right| \lesssim U_n^{-(s+2)},$$

$$\left| \int_0^{\infty} w_{\lambda}^{U_n}(u) \operatorname{Re}(\mathcal{F}v(u)) \, du \right| \lesssim U_n^{-(s+1)}.$$

**Ex. 10.2.** Denote by  $\mathcal{F}f$  the Fourier transform of  $f$ . For an integer  $s \geq 1$  suppose that  $v$  satisfies  $\max_{k \in \{0, 1, \dots, s\}} \|v^{(k)}\|_{L^2(\mathbb{R})} \leq C$  and  $\|v^{(s)}\|_{\infty} \leq C$  for some  $C > 0$ , where  $v^{(k)}$  is the  $k$ -th derivative of  $v$ . Let  $w$  be a function supported on  $[-1, 1]$  such that  $(1 - w(\bullet))/(\bullet)^s \in L^2(\mathbb{R})$  and  $\mathcal{F}[(1 - w(\bullet))/(\bullet)^s] \in L^1(\mathbb{R})$ . Show that

$$\|\mathcal{F}^{-1}[(1 - w(\bullet/U_n))\mathcal{F}v(\bullet)]\|_{\infty} \lesssim U_n^{-s}.$$

**STATISTICS FOR STOCHASTIC PROCESSES - EXERCISE SHEET 11**  
**(not part of the course in 2024)**

Mastermath, Spring Semester 2024, Jakob Söhl (j.soehl@tudelft.nl)

**Ex. 11.1.** Let  $X_1, X_2, \dots, X_n$  i.i.d. random variables. Let  $\varphi_n$  be the empirical characteristic function of  $X_1, X_2, \dots, X_n$  and  $\varphi_n^{(k)}$  the  $k$ -th derivative of  $\varphi_n$ . For complex-valued random variables  $Z$  we define  $\text{Var}_{\mathbb{C}}(Z) = \mathbb{E}[|Z - \mathbb{E}[Z]|^2]$ . Show that for  $k = 0, 1, 2$  and for all  $u \in \mathbb{R}$

$$\text{Var}_{\mathbb{C}}\left(\varphi_n^{(k)}(u)\right) \leq \frac{1}{n} \mathbb{E}[X_1^{2k}]$$

## STATISTICS FOR STOCHASTIC PROCESSES - EXERCISE SHEET 12

Mastermath, Spring Semester 2024, Jakob Söhl (j.sohl@tudelft.nl)

**Ex. 12.1.** Let  $X$  be a one-dimensional Lévy process with Lévy measure  $\nu$  satisfying  $\int_{\mathbb{R}} x^2 d\nu(x) < \infty$ . Let  $\mathbb{P}_{\Delta}$  be the probability measure of  $X_{\Delta}$ . Show that for every bounded continuous function  $f : \mathbb{R} \rightarrow \mathbb{R}$

$$\int_{\mathbb{R}} f(x)x^2 \frac{d\mathbb{P}_{\Delta}(x)}{\Delta} \rightarrow \sigma^2 f(0) + \int_{\mathbb{R}} f(x)x^2 d\nu(x) \quad \text{as } \Delta \rightarrow 0.$$

[You may use the following fact: For a sequence of probability measures  $\mathbb{P}, \mathbb{P}_1, \mathbb{P}_2 \dots$  on  $\mathbb{R}$  with characteristic functions  $\varphi, \varphi_1, \varphi_2, \dots$  a consequence of Lévy's continuity theorem is that the pointwise convergence  $\varphi_n \rightarrow \varphi$  as  $n \rightarrow \infty$  implies that  $\mathbb{P}_n$  converge weakly to  $\mathbb{P}$  as  $n \rightarrow \infty$ .]

**Ex. 12.2.** Let  $X$  and  $\epsilon$  be independent real-valued random variables whose distributions are absolutely continuous with respect to the Lebesgue measure with Lebesgue densities  $p_X$  and  $p_{\epsilon}$ , respectively. Let  $\varphi_{\epsilon}$  and  $\varphi_Y$  be the characteristic functions of  $\epsilon$  and  $Y = X + \epsilon$ , respectively. Let  $\varphi_{Y,n}$  be the empirical characteristic function of  $n$  i.i.d. copies of  $Y$ . Suppose  $\varphi_{\epsilon} \neq 0$  for all  $u \in \mathbb{R}$  and let  $M_U := \sup_{|u| \leq U} |1/\varphi_{\epsilon}|$ . Assume for an integer  $s \geq 1$  that  $p_X^{(k)} \in L^2(\mathbb{R})$  for all  $k \in \{0, 1, \dots, s\}$ . Define

$$\hat{p}_X = \mathcal{F}^{-1} \left[ \frac{\varphi_{Y,n}}{\varphi_{\epsilon}} \mathbb{1}_{[-U_n, U_n]} \right].$$

Show that

$$\|\hat{p}_X - p_X\|_{L^2(\mathbb{R})} = O_{\mathbb{P}} \left( U_n^{-s} + M_{U_n} U_n^{1/2} n^{-1/2} \right).$$

## STATISTICS FOR STOCHASTIC PROCESSES - EXERCISE SHEET 13

Mastermath, Spring Semester 2024, Jakob Söhl (j.soehl@tudelft.nl)

**Ex. 13.1.** Let  $N = (N_t)_{t \geq 0}$  be a Poisson process with intensity  $\lambda \geq 0$ . For  $\Delta > 0$  we define  $Z_i := N_{i\Delta} - N_{(i-1)\Delta}$  with  $i = 1, 2, \dots, n$ .

a) Compute the maximum likelihood estimator of  $\lambda$  given the observations  $\mathbb{1}_{Z_i \neq 0}$ ,  $i = 1, \dots, n$ . Compare to the nonlinear estimator  $\tilde{\lambda}_n$  from the lecture.

b) Let

$$Y_i := \begin{cases} 0 & \text{if } Z_i = 0, \\ 1 & \text{if } Z_i = 1, \\ 2 & \text{if } Z_i \geq 2, \end{cases}$$

for  $i = 1, \dots, n$ .

Find the log-likelihood for estimating  $\lambda > 0$  from the observations  $Y_1, Y_2, \dots, Y_n$ . Set its derivative equal to zero and approximate the solution of the resulting equation for high-frequency observations  $\Delta \rightarrow 0$  as  $n \rightarrow \infty$ . Compare the resulting estimator to the linear estimator  $\hat{\lambda}_n$  from the lecture.

## STATISTICS FOR STOCHASTIC PROCESSES - EXERCISE SHEET 14

Mastermath, Spring Semester 2024, Jakob Söhl (j.soehl@tudelft.nl)

**Ex. 14.1.** Suppose that  $(X_i)_{i=1}^n$  are zero-mean and independent random variables such that, for some fixed  $q \geq 1$ , they satisfy the moment bound  $(\mathbb{E}[|X_i|^{2q}])^{\frac{1}{2q}} \leq K_q$ . Show that

$$\mathbb{P} \left( \left| \frac{1}{n} \sum_{i=1}^n X_i \right| \geq \delta \right) \leq B_q \left( \frac{1}{\sqrt{n}\delta} \right)^{2q} \quad \text{for all } \delta > 0,$$

where  $B_q$  is a universal constant depending only on  $K_q$  and  $q$ . [Hint: You may use Rosenthal's inequality.]

**Ex. 14.2.** Let  $X$  be a real-valued Lévy processes. Let  $D := [a, b] \subseteq \mathbb{R} \setminus \{0\}$  and

$$\mathcal{S} := \{\beta_1 \varphi_1 + \dots + \beta_d \varphi_d \mid \beta_1, \dots, \beta_d \in \mathbb{R}\},$$

where  $\varphi_j$  with  $j = 1, \dots, d$  are  $\nu$ -a.e. continuous, bounded functions which have support in  $D$  and are orthonormal with respect to the inner product  $\langle p, q \rangle := \int_D p(x)q(x) dx$ . Let  $0 = t_0 < t_1 < \dots < t_n$  and define

$$\hat{\rho}(x) := \sum_{j=1}^d \hat{\beta}(\varphi_j) \varphi_j(x) \quad \text{with} \quad \hat{\beta}(\varphi_j) := \frac{1}{t_n} \sum_{k=1}^n \varphi_j(X_{t_k} - X_{t_{k-1}}).$$

Show that  $\hat{\rho}$  is the unique solution of the minimisation problem

$$\min_{f \in \mathcal{S}} \gamma_D(f),$$

where  $\gamma_D : L^2(D, dx) \rightarrow \mathbb{R}$  is given by

$$\gamma_D(f) := -\frac{2}{t_n} \sum_{k=1}^n f(X_{t_k} - X_{t_{k-1}}) + \int_D f^2(x) dx.$$

## STATISTICS FOR STOCHASTIC PROCESSES - EXERCISE SHEET 15

Mastermath, Spring Semester 2024, Jakob Söhl (j.soehl@tudelft.nl)

**Ex. 15.1.** Let  $X$  be a one-dimensional Lévy process with Lévy measure  $\nu$ . Let  $\varphi$  have support in  $[c, d] \subseteq \mathbb{R}_{>0}$  and let  $\varphi|_{[c,d]}$  be continuous with continuous derivative.

a) Show that

$$\begin{aligned}\mathbb{E}[\varphi(X_\Delta)] &= \varphi(c) \mathbb{P}(X_\Delta \geq c) - \varphi(d) \mathbb{P}(X_\Delta > d) + \int_c^\infty \varphi'(u) \mathbb{P}(X_\Delta \geq u) \, du, \\ \int_{-\infty}^\infty \varphi(x) \, d\nu(x) &= \varphi(c) \nu([c, \infty)) - \varphi(d) \nu((d, \infty)) + \int_c^\infty \varphi'(u) \nu([u, \infty)) \, du.\end{aligned}$$

b) Derive further

$$\left| \frac{\mathbb{E}[\varphi(X_\Delta)]}{\Delta} - \int_{-\infty}^\infty \varphi(x) \, d\nu(x) \right| \leq \left( |\varphi(c)| + |\varphi(d)| + \int_c^d |\varphi'(u)| \, du \right) M_\Delta([c, d]),$$

where  $M_\Delta([c, d]) := \lim_{\epsilon \rightarrow 0} \sup_{y \in [c, d+\epsilon]} \left| \frac{1}{\Delta} \mathbb{P}(X_\Delta \geq y) - \nu([y, \infty)) \right|$ .